

Patient Awareness

Disease Process	Why it Matters	Who's at Risk or Patients with:	Signs and Symptoms	Diagnostic
Stroke	Third leading cause of death in the US. #1 reason for long term disability.	 High Cholesterol High Blood Pressure Diabetes Smokers	 Confusion / Memory changes Trouble speaking One sided weakness Vision disturbance Dizziness, passing out, feeling faint 	Carotid Ultrasound to look for plaque in the arteries, leading to more risk for TIA or stroke.
Heart Failure	 #1 Cause of death in Americans. 60% of deaths are from Sudden Cardiac Death. 	High Blood PressureHigh CholesterolDiabetesCADValve DiseaseSmokers	Chest discomfort, pressure or pain Shortness of breath Leg heaviness or swelling Palpitations or flutter feeling in chest	Echocardiogram to evaluate the heart muscle function, size and valve function.
Vascular Disease	 PAD affects 8 to 12 million Americans each year. Narrowing of blood vessels that carry blood to the legs. 	Diabetics Smokers CAD	Leg / arm swelling Leg / arm Heaviness Leg / arm discomfort / pain Buttocks pain Leg/arm coldness	Arterial Doppler, Venous Doppler, and ABI to evaluate for plaque ulceration or thrombus/DVT in legs or arms.
Aorta (Abdominal)	 AAA is known as the silent killer. A ruptured aortic aneurysm can rapidly lead to life threatening internal bleeding. 	SmokersArtherosclerosisFamily HistoryHigh Blood PressureTrauma	Back or Stomach Pain Tingling / numb legs / feet Cold legs / feet Swelling in legs	Aortic Ultrasound to evaluate for plaque, dilatations, and abdominal aortic aneurysm.
Abdominal	Abdominal diseases affects nearly 20% of Americans.	Smokers Obese, High BMI Abnormal labs	Abdominal painNausea / vomitingDiarrheaGas / bloating	Abdominal Ultrasound to evaluate organs in the abdomen.
Kidney Disease	• 9th leading cause of death in the United States.	DiabetesHigh Blood PressureBlood in urineSmokers	Swelling of extremitiesBlood in urineDecreased urine output	Renal Ultrasound with Bladder and Renal Artery to evaluate function.
Thyroid	Thyroid Disease and thyroid lesions are very common and in most cases benign.	Men and Women Family history Lumps in neck	Fatigue Abnormal labs Unusual weight gain	Thyroid Ultrasound to evaluate for abnormalities.
Neuropathy	 25-30% of population will be affected by neuropathy. 60-70% of people with diabetes have neuropathy. 	Diabetes Carpel Tunnel Trauma	Extremity discomfortExtremity painExtremity numbnessExtremity tingling	Nerve Conduction Study to evaluate nerves.

Incontinence Or Weak Bladder	 Millions of men and women around the country suffer from incontinence. Incontinence can be reduced or eliminated with pelvic therapy. 	Women with multiple child births Men following prostate surgery Age	· Unintentional urinary release or leakage from sneezing, coughing, laughing, exercise, or other activities.	Urodynamic flow study to evaluate bladder function. PFR therapy to eliminate or reduce incontinence.
------------------------------------	--	---	--	---